

# Brisbane South Antenatal Shared Care

## Process

### First GP Visit(s)

(May take more than one consultation)

- Confirm pregnancy & dates. Scan after 7/40
- Scan if dates uncertain **or** risk of ectopic (previous ectopic, tubal surgery) **or** previous pregnancy complications/medical risks
- Folate and iodine supplementation for all
- Review medical, surgical, psych, family history, medications, allergies etc.- update GP records ± create My Health Record shared health summary.
- Identify risk factors for pregnancy.
- Discuss and offer genetic carrier testing, anomaly screening +/- NIPT.
- BP, weigh, calculate BMI, physical examination.
- Discuss smoking, nutrition, alcohol, physical activity; dietary advice (listeria) & drug avoidance; Assess emotional well-being and screen for DFV if safe to do so.
- Consider early Aspirin use if risk factors for pre-eclampsia/IUGR – before 16 weeks (see over)
- Offer influenza and COVID (follow current guidelines) vaccination as soon as practical.
- Discuss models of care

### First Trimester Screening Tests

(cc. ANC on all request forms) – all requested tests to be reviewed and actioned by referring clinician.

- FBC, Ferritin, blood group and antibodies, rubella, Hep BsAg, Hep C, HIV, syphilis serology, MSU (treat asymptomatic bacteriuria)
- Discuss and offer Genetic Carrier Screening to ALL
- Discuss and offer screening for anomalies in ALL:
  1. Nuchal Translucency Scan + First Trimester Screen (free  $\beta$ hCG, PAPP) K11-13<sup>+</sup>6 **OR**
  2. Non-Invasive Prenatal Testing > K9 (Higher failure rate in multiple pregnancy, not Medicare funded, first trimester scan recommended) **OR**
  3. Triple Test (AFP, Oestriol, hCG) K15-22 if too late for first TM testing. Not if twins or diabetes. Discuss/ offer CVS/Amniocentesis if appropriate.
- Order Rhesus-D NIPT of Rh Negative non-alloimmunised patients (timing is lab dependent.)
- Cervical screening test if due, Dry swab (PCR) if lesions/chancres present.
- Varicella serology (if no varicella history /vaccination)
- OGTT after K12 (or HbA1c) if high risk for Diabetes (see box below)
- ELFT, TFTs, Vit D, chlamydia/gonorrhoea **only recommended for at risk women (see over)**

### Uncomplicated pregnancy

- Refer privately for detailed scan (placenta, morphology, cervical length) at 18-20 weeks.
- First Midwifery Booking visit at 14-16/40 with medical visit at 14-20/40 (18-20/40 combined RM/doctor visit MMH)
- **You are responsible for her care until she is seen by the hospital, after which the responsibility is shared.**
- GP visits to be scheduled around hospital appointments to ensure timely review of results.
- **All investigations to be reviewed by referring clinician and required follow up arranged or referrals made.**

### GP Visits: 14, 24, 28, 31, 34, 38, 40 weeks

(More frequent if clinically indicated)

- Record or place printed copy of notes/ results in Pregnancy Health Record (PHR)
- Schedule, education, and assessment as per the PHR
- K26-28 GTT, FBC, Ferritin, Blood group and antibody screen, Syphilis Serology
- Syphilis PCR (dry swab) anytime as clinically indicated).
- K36 Hb, (Ferritin if indicated), Syphilis serology)
- **Vaccinations:** Offer influenza & COVID (any time); pertussis at K20-32 in each pregnancy & Abrisvo (RSV) between 28-36 weeks gestation.
- **ANC review** at K36 and at K40-41

### Pre-Conception Unique role for GPs!

- Folate and iodine supplementation for all
- Rubella serology +/- vaccination
- Varicella serology if no history +/- vaccination
- Influenza Vaccination in season + and COVID (follow current guidelines)
- Cervical screening if due
- Chlamydia test/treat <30yrs
- Smoking cessation
- Alcohol cessation
- Discuss and offer reproductive carrier screening e.g., CF, SMA & FXS (or extended panel)
- Consider referral to preconception clinic e.g., Mater, Logan Pre-pregnancy assessment.

### Rh Neg Mothers with unknown or positive fetal Rh status

- Antibody negative. offer 625 IU anti-D at 28 & 34 weeks' and for sensitising events.
- Dose can be given at local Hospital, OR GP—order via QML or Mater Blood Bank, delivered via courier to surgery.
- QML 3371 9029
- Mater 3163 8179
- AntiD not indicated for threatened miscarriage  $\leq$  12/40 (or ToP  $\leq$  10/40)

### High Diabetes in Pregnancy Risk Please specify reason and include copy of results in referral.

- Previous GDM or baby > 4500g, PCOS, strong family hx, BMI > 30, maternal age  $\geq$  40, previous unexplained perinatal loss; multiple preg, high risk ethnicity, glycosuria; assisted reproduction; Medications – steroids, antipsychotics
- HbA1c up to 12/40 if early screening indicated, Consider OGTT at >10/40 as clinically indicated & patient tolerated. Avoid OGTT in post bariatric surgery patients.
- **URGENT Hospital ANC referral if – GDM = HbA1c 6-6.4%; Fasting  $\geq$  5.3 -6.9 mmol or 1-hr  $\geq$  10.6 or 2-hr 9-11 OR OVERT DIP = HbA1c  $\geq$  6.5%; Fasting  $\geq$  7.0 mmol or 2-hr  $\geq$  11.1 mmol**

### Medical or Obstetric Complications? EARLY or URGENT ANC referral:

- GP referral letters are triaged by MW or consultant within same week.
- Please specify urgency and reasons in the referral letter
- Refer to local service – will liaise or make further referrals if required.
- **Be sure to cc pathology and radiology and give women a copy of their results.**
- Cervical length < 35mm transabdo USS – arrange TVS; If < 25mm (TVS) commence 200mg vaginal progesterone daily; If < 10mm, URGENT referral? cerclage

| CONTACTS   | Beaudesert                                     | Logan  | Redland                           | Mater  |
|--|--|--|-----------------------------------|--|
| Secure e-Referral  | SMART Referrals or Medical Objects/Health Link |  |                                   |  |
|  | Central Referral Hub: 1300 364 248             |  |                                   | 3163 8053                                      |
| Updated information to be sent via Smart Referral (or ANC Fax)   | 5541 9132                                      | 2891 6976  | 3488 3436                         | 3163 8053                                      |
| ANC phone  | 5541 9144                                      | 3086 3379  | 3488 3434                         | 3163 1861                                      |
| Perinatal Mental Health Services   | 3089 2734                                      | 3089 2734  | 3825 6214                         | 3163 7990                                      |
| GP Liaison Midwife   | 0482 677 946 or GPLO GP- 2891 5754             |  |                                   | 3163 1861                                      |
| <b>For Urgent Referral or Advice</b>   |  |  |                                   |  |
| O&G Registrar  | -  | 2891 8027  | 3488 3758                         | 3163 6611                                      |
| Obstetrician/GP Obs on call  | 5541 9174                                      | 3089 6963  | 3488 3111                         | 3163 6612                                      |
| Triage Midwife   | 5541 9181                                      | 2891 8811  | 3488 3044                         | 3163 1861                                      |
| <b>For urgent MH referral/advice</b>   | 1300 642255 (1300 MHCALL) for all centres      |  |                                   |  |
| <b>Pregnancy Complications</b>   |  |  |                                   |  |
| Complications e.g., bleeding, pain, incomplete miscarriages, altered fetal movts. (Logan EPAU - business hours only)<br><b>Haemodynamically unstable women? Direct to ED/PAC</b> | On-Call GP Obstetrician<br>5541 9174           | <14w 2891 8456<br>>16w Phone MAC Reg –<br>2891 8027<br>ED: 2891 8899 | On-Call Obstetrician<br>3488 3111 | Pregnancy Assessment Centre (PAC)<br>3163 6577 |

# Maternity GP Shared Care

## Additional Information and Advice

### Additional Tests – STI screen, ELFT, TSH/TFTs, Vit D, TORCH serology

- Chlamydia and Gonorrhoea –test women < 30 years old and other high-risk women by self-collect PCR swab.
- ELFTs and urinary protein/creatinine ratio recommended for obese women (BMI > 30), hypertension or known or suspected renal or liver disease, autoimmune disease.
- Routine TFTs *are not* recommended in low-risk pregnant women. TSH generally drops in first trimester with the rise in HCG. If a woman has a TSH lower than the lab reference range, check free T4/T3—if these are normal, the woman *does not* need referral, if elevated, they will need clinical review, possibly referral – liaise with your local team. <https://metronorth.health.qld.gov.au/wp-content/uploads/2017/10/thyroid-disorders-pregnancy.pdf>
- Women with pre-existing hypothyroidism should have a TSH <2.5 in first trimester and <3.0 in the rest of the pregnancy. Lab reference ranges will reflect pregnancy recommendations if the woman is identified as being pregnant. Weekly doses usually need to increase by 30% during pregnancy, which is an extra 2 doses/week. Advise women to commence the higher dose as soon as they know they are pregnant.
- Vitamin D levels or supplementation are recommended for obese or dark-skinned women or those with little sun exposure or who cover themselves for religious or cultural reasons. Levels <50 may require supplements of 2000 IU/day. Levels <15 require higher doses and re-test after 3 months.
- Toxoplasma, cytomegalovirus, and herpes serology should *not* be performed routinely. If risk factors indicate a need for testing, please include risk in your referral as follow-up tests or other investigations or management may be needed.

### Nutrition and Supplements

- Folate - 0.5 mg for all low risk, 2.5-5 mg if high risk (diabetic, obese, previous, or familial neural tube defect, anticonvulsants). Start one month before conception & continue to 12 weeks.
- Iodine 150mcg/day - recommended preconception, during pregnancy and while breastfeeding (folate + iodine supplement is available)
- 2-3 serves daily of calcium-rich food/drink (1g/day) OR add 500mg minimum daily supplement. RANZCOG recommend universal 400IU/day Vitamin D (e.g., 600mg Ca + 1000IU Vit D)
- Iron only needed if deficiency is identified however low dose is included in all pregnancy supplements. Avoid Vit A in pregnancy.
- Added supplements needed for women post Bariatric Surgery (including Vitamin A) – seek Dietitian input.
- Avoid or limit intake of large/predatory fish due to mercury content (Orange Roughy /Sea Perch, Shark/Flake, Swordfish, Marlin etc.)

### Preventing Infections

- Toxoplasmosis - Avoid feeding raw/undercooked meats to pets, avoid cat faeces/litter, wear gloves when gardening.
- Cytomegalovirus - Good hand hygiene; Care with urine, saliva, nappies of young children
- Influenza and COVID Vaccination at any stage antenatally; pertussis vaccinations 20-32 weeks & RSV 28-36 weeks (but up to time of delivery if missed; requires two weeks to be fully effective)
- Listeriosis - Avoid soft cheeses, un-pasteurised milk, pate, raw eggs, hot dogs, undercooked and deli meats, reheated leftovers, precut fruit, bean sprouts.

### Early Low Dose Aspirin (100-150mg)

Commence before 16/40 (stop at 36/40) to reduce incidence of placental disorders such as Pre-eclampsia & fetal growth restriction (FGR), preterm birth & perinatal mortality in those at increased risk. Take in the evening.

#### High Risk Factors - recommend if patient has one or more of:

- Hypertension
- Renal disease
- Auto-immune diseases e.g., SLE or anti-phospholipid syndrome
- Diabetes (Type 1 or Type 2)
- Previous History of pre-eclampsia

#### Moderate Risk Factors – consider if two or more are present:

- Primiparous
- BMI > 35
- Age > 40
- Multiple pregnancy
- Family history of pre-eclampsia (mother or sister)
- More than 10 years since last pregnancy

### More Online Information and Education for GPs interested in Antenatal Care are available through:

- General Practice Liaison Officer (GPLO) Program webpage: <https://metrosouth.health.qld.gov.au/referrals/general-practice-liaison-officer-gplo-program>
- Mater Mothers [www.materonline.org.au](http://www.materonline.org.au) (Click on Shared Care Alignment for a range of resources for GPs) [www.matermothers.org.au](http://www.matermothers.org.au) (Click on Mater Mothers' Hospital for resources for women)
- [www.maternity-matters.com.au](http://www.maternity-matters.com.au) has consumer and clinician resources and links to reputable websites.

### Early Pregnancy Complications (<20 weeks)

- Nausea and vomiting - decrease iron (but continue iodine and folate), try ginger, acupuncture, pyridoxine 75 mg/day in divided doses, doxylamine (Cat A) Metoclopramide (Maxolon Cat A) and Phenothiazines like Prochlorperazine (Stemetil Cat C, po/pr/iv, safe in first trimester); Ondansetron may be effective but is relatively expensive. Even mild dehydration/ketonuria may benefit from IV fluids.
- Bleeding: check blood group and antibodies. Threatened miscarriage in rhesus-negative women without antibodies (and unknown fetal Rh status) after 12 weeks requires anti-D, before 12 weeks anti-D is not required unless the miscarriage completes, or you are concerned the woman may not re-present.
- Bleeding and pain: consider ectopic pregnancy!
- Consider advice from, or referral to, early pregnancy assessment unit (EPAU), pregnancy assessment centre (PAC) or emergency department at booking hospital (appointments may be required)

**Beaudesert 5541 9111; Logan EPAU (< K14) 3299 8456 – 8-4 Mon-Fri only  
Redlands 3488 3111; Mater PAC 3163 6577**

### Late pregnancy complications (>16/40 at Logan; >20/40 at other Brisbane South maternity hospitals)

- Bleeding – can do spec exam but avoid PVE. Exclude cervical dilation. Re-check placental site on original morphology scan, Rhesus neg mothers (with unknown fetal Rh Status) need anti-D.
- Abdominal pain - can do spec exam but no PVE. Exclude cervical dilation.
- Ruptured membranes - Review at hospital preferred. Can do spec exam but no PVE.
- Fundal height > 3cm above or below expected for gestational age – arrange USS & if IUGR confirmed, refer to ANC by Fax *and* Phone Obstetrician/Registrar; if LGA confirmed, refer back through ANC
- Perceived change in fetal movements beyond 28 weeks or no FH detected – arrange IMMEDIATE hospital review.
- Most should be referred to booking hospital birth suites, pregnancy/maternity assessment/observation units or Emerg. Dept.

**Beaudesert 5541 9111; Logan MAC  
Redlands 3488 3111; Mater PAC 3163 6577**

For feedback on this document, please email MSHHS GPLO Maternity Team at [GPLO\\_Maternity\\_Share\\_Care@health.qld.gov.au](mailto:GPLO_Maternity_Share_Care@health.qld.gov.au)